

Optics, Spring 2016

Exercise 6, 1.3.2016

1. Field of view (Ch 5.7)

A rectangular CCD sensor (22.2 mm x 14.8 mm) is placed to the focal plane of the objective lens ($f = 1000$ mm).

- Calculate the field of view (FOV) of the arrangement, i.e. the angular diameter of the object at infinity, which corresponds to the CCD sensor size.
- Is the FOV sufficient for the photography of the Moon (30') or the Andromeda galaxy (90' x 100')?

2. Chromatic aberration (Ch. 5.2)

Due to chromatic aberration, the focal length of a lens depends on wavelength. Assume that focal lengths f_1 and f_2 correspond to refractive indices n_1 and n_2 respectively. Show that, if $n_3 = \frac{n_1+n_2}{2}$ corresponds to focal length $f_3^2 \approx f_1 f_2$, the difference in focal lengths is $f_1 - f_2 = \frac{n_2-n_1}{n_3-1} f_3$.

3. Spherical mirrors (Ch 5.4)

- A 1-cm tall object is positioned 12 cm in front of a spherical concave mirror having a radius of curvature of 8 cm. Calculate the image location, size, orientation and type. Draw a ray diagram.
- Repeat the calculation for a convex mirror with $f = 9$ cm. Draw a ray diagram.

4. Mirror magnification (Ch 5.4)

Derive the expression for the transverse magnification of a spherical mirror (concave and convex) in paraxial approximation.

5. Spherical mirrors continued (Ch 5.4)

The imaging equation for large spherical mirrors is:

$$\frac{1}{s_0} + \frac{1}{s_i} = -\frac{2}{R} - \frac{H^2}{R} \left(\frac{1}{R} - \frac{1}{s_0} \right)^2$$

where H is the height on the mirror (see the figure). Estimate the amount of spherical aberration by calculating the difference between focal lengths $f - f'$ when the object is in infinity and $H \ll R$. Here f and f' correspond to a paraxial ray (parallel to optical axis) and a ray reflecting from the top part of the mirror, respectively. Calculate the difference when $H = 8$ cm and $R = 10$ cm.

